

Flux-Bubble Models and Mesonic Molecules

M.M. Boyce^a, J. Treurniet, and P.J.S. Watson

*Ottawa-Carleton Institute for Physics, Carleton University, Ottawa, Ontario,
Canada, K1S-5B6*

It has been shown that the string-flip potential model reproduces most of the bulk properties of nuclear matter, with the exception of nuclear binding. Furthermore, it was postulated that this model, with the inclusion of the colour-hyperfine interaction, should produce binding. In some recent work a modified version of the string-flip potential model was developed, called the flux-bubble model, which would allow for the addition of perturbative QCD interactions. In attempts to construct a simple $q\bar{q}$ nucleon system using the flux-bubble model (which only included colour-Coulomb interactions) difficulties arose with trying to construct a many-body variational wave function that would take into account the locality of the flux-bubble interactions. In this talk we consider a toy system, a mesonic molecule in order to understand these difficulties. *En route*, a new variational wave function is proposed that may have a significant enough impact on the old string-flip potential model results that the inclusion of perturbative effects may not be needed.

1 Introduction

For the past 30 years several attempts have been made, with little success, to describe nuclear matter in terms of its constituent quarks. The main difficulty is the non-perturbative nature of QCD. The only rigorous method for handling multi-quark systems to date is lattice QCD. However, this is very computationally intensive and given the magnitude of the problem it appears unlikely to be useful in the near future. As a result, more phenomenological means must be considered.

A good phenomenological model should be able to reproduce, at least qualitatively, all the overall bulk properties of nuclear matter, in particular:

- nucleon gas at low densities with no power-law van der Waals forces
- nucleon binding at higher densities
- nucleon swelling and saturation of nuclear forces with increasing density
- quark gas at extremely high densities.

There are many models that attempt to reproduce these properties but none of them does so completely. In this talk, only the string-flip potential model^{1,2,3} will be considered. This model appears to be promising because it reproduces most of the aforementioned properties with the exception of nucleon binding.

^aSpeaker.

In an attempt to find the root of this problem, a toy mesonic-molecular system will be used to study the inclusion of perturbative interactions and variational wave functions.⁴

This proceeding is essentially part II of our MRST '94 talk.⁵ For completeness sake, a brief overview of the previous proceedings will be given, followed by our most recent work.

2 From String-Flips to Flux-Bubbles

In this a section, a brief outline of the generic string-flip potential model is presented, followed by an overview of the flux-bubble model.

2.1 The String-Flip Potential Model

The string-flip potential model postulates how flux-tubes should form between quarks at zero temperature, based on input from lattice QCD and experiment. An adiabatic assumption is made, in which the quarks move slowly enough for their fields to reconfigure themselves, such that the overall potential energy is minimized: *i.e.*,

$$V = \sum_{\min\{q_m \dots q_n\}} \mathbf{v}(\vec{r}_m \dots \vec{r}_n) = \left\{ \begin{array}{c} \text{Diagram of a cube containing a complex network of lines (flux-tubes) connecting various points (quarks) inside and on the surface of the cube.} \end{array} \right\}. \quad (1)$$

In Eq. 1, the sum is over all gauge invariant sets, $\{q_m \dots q_n\}$, of N_q quarks, $\{q_1 \dots q_{N_q}\}$, in a box of side L with periodic boundary conditions (to simulate continuous nuclear matter).

The physics, or model input, is contained within the many-body terms, $\mathbf{v}(\vec{r}_m \dots \vec{r}_n)$, in V , which, in general, can be quite complex and virtually impossible to compute. Ironically, the form of $\mathbf{v}$ is quite simple: the total minimal length, ℓ , of flux-tubing (or string) connecting a given set of quarks times the string tension, σ , $\mathbf{v} \sim \sigma \ell$. Over the past couple of years, many simplifications of $\mathbf{v}$ have been assumed in the form of two-body, three-body, and long complex chains of quarks, using $SU_c(2)$ and $SU_c(3)$ with and without “moving-colour.”^{b1,2,3} All of these models yield similar results for nuclear matter —

^bTypically, a quark is assigned a “fixed-colour” which doesn’t change throughout the evolution of the gas. Conversely, “moving-colour,” is when the quarks are allowed to change colour.

no nuclear binding. A general synopsis of these models can be found in the MRST '94 proceedings.⁵ For simplicity, we shall concentrate on the two-body quark-antiquark form, $\mathbf{v} \sim \sigma r_{q\bar{q}}$, with $SU_c(2)$.

2.2 The Flux-Bubble Model


To date, string-flip potential models assumes only non-perturbative interactions: *cf.*, $\mathbf{v} \sim \sigma r - \alpha_s/r$. Flub-bubble models are a recent attempt to include perturbative interactions. In particular, they are a derivative of the following two-body potential⁵

$$\mathbf{v}_{ij} = \sigma(r_{ij} - r_0)\theta(r_{ij} - r_0) + \alpha_s \lambda_{ij} \left(\frac{1}{r_{ij}} - \frac{1}{r_0} \right) \theta(r_0 - r_{ij}), \quad (2)$$

where λ_{ij} is a model dependent colour factor, which is free from Van der Waals forces.

A “flux-bubble” is formed when two or more ends of neighboring flux-tubes come within r_0 of each other, by inserting virtual $q\bar{q}$ -pairs on the boundary of this region, thus leaving a perturbative bubble with non-perturbative flux-tubes emanating from it.

E.g. Consider configuration (a) of quarks, with $r > r_0$, about to move to (b), *s.t.*, two of them are within $r < r_0$. Then the procedure is to draw a bubble of r_0 away from the two, (b), and to cut the flux-tubes at the boundary and insert virtual $q\bar{q}$ pairs, (c). Once the potential is computed the configuration is restored to (b) before the next move is made.



Note the insertion of the virtual $q\bar{q}$ pairs allows the construction of colourless objects. These are solely used as a tool to calculate the overall length of the flux-tube correctly, and not used in computing the Coulomb term, however, as the field energy is already taken into account by the “real” quarks inside the bubbles. In general, once the bubbles have been determined, the flux-tubes must be reconfigured in order to minimize the linear part of the potential.

Therefore, the most general $q\bar{q}$ -flux-bubble potential, with $SU(2)$ -colour, is given by (*cf.*, Eq. 1)

$$V = \sigma \sum_{\min\{q\bar{q}\}} (r_{q\bar{q}} - r_0) \theta(r_{q\bar{q}} - r_0) + \alpha_s \sum_{i < j} \lambda_{ij} \left(\frac{1}{r_{ij}} - \frac{1}{r_0} \right) \theta(r_0 - r_{ij}), \quad (3)$$

where λ_{ij} equals $-3/4$ for unlike-quarks and $1/4$ for like-quarks. The first term represents the string-flip, or linear, potential model, while second term contains Coulomb corrections to the this potential (*i.e.*, linear-plus-Coulomb potential model) plus new terms describing interactions between the ends of the flux-tubes.^c

2.3 Implementation Problems

The actual implementation of the flux-bubble potential turned out to be quite straightforward. However, trying to find a good many-body variational wave function proved to be insurmountable. The main problem was that each trial wave function required vast amounts of calculations by hand, writing computer code, and lots of computer time (running several machines in tandem). Human CPU time ($\tau_{\text{human}} \sim \mathcal{O}(1)wk$) aside, using a simple mesh minimization scheme requires $\mathcal{O}(M^p/m)$ of CPU time (τ_{CPU}), where M is the mesh size, p is the number of parameters, and m is the number of machines.

A string-flip potential model, for 7 triplets of quarks using fixed $SU(2)$ -colour, with a 3-parameter wave function,

$$\Psi_{\alpha\beta} = e^{\underbrace{- \sum_{\min\{q\bar{q}\}} (\beta r_{q\bar{q}})^\alpha}_{\text{Correlation} \Leftrightarrow \beta}} \underbrace{\prod_{\text{colour}} |\Phi(r_{p_k})|}_{\text{Slater} \Leftrightarrow \rho}, \quad (4)$$

with parameters ρ (density), β (inverse correlation length), and α , would consume about 5 weeks of CPU time on our 8-node farm, here at Carleton, using a crude 10^3 mesh. However, because of certain assumptions ($\alpha(\rho) \approx \alpha(\rho = 0)$) and scaling tricks ($(\beta, \rho^{1/3}) \rightarrow \zeta(\theta) (\cos \theta, \sin \theta)$), the CPU time can be reduced to 8hrs. Unfortunately, because of the local nature of the flux-bubble model, we cannot use scaling tricks. So, in general, we would require $\tau_{\text{CPU}} \sim (10^{p-1}/21)wks!$

Thus, a way of checking different wave functions and minimization schemes which do not consume large amounts of CPU time is desirable. In particular,

^cIt is conceivable, that by extending the model to include interactions along the length of the flux-tubes, could lead to a model for pion-liquid crystals.⁶

a mini-laboratory is needed in which various aspects of the string-flip and flux-bubble potential models, from wave functions to minimization schemes, can be investigated, without being too concerned about *CPU* overhead.^d

3 Mesonic Molecules

An interesting place that might make a good mini-laboratory is a mesonic-molecule,⁷ Q_2 , consisting of two heavy-quarks, Q , and two relatively light antiquarks, $\bar{q}$. The heavy-quarks are assumed to be massive enough so as not to be disturbed by the motion of the light-antiquarks: *i.e.*, the adiabatic approximation. By varying the distance, R , between the heavy-quarks a mesonic-molecular potential, $U(R)$, can be computed.

The Schrödinger equation that describes the effective potential, $U(R)$, is given by⁸

$$\left(\frac{1}{2m_q} \sum_{\bar{q}} \vec{\nabla}_{\bar{q}}^2 + V \right) \Psi = U(R) \Psi \quad \text{---} \quad \begin{array}{c} \bar{q} \circ \\ \swarrow \\ \bullet_q \text{---} R \text{---} \bullet_q \\ \searrow \\ \circ_{\bar{q}} \end{array} \quad (5)$$

where $m_q = 330 \text{ MeV}$. The potential, V , describes the many-body nature of the four quark system and is therefore model dependent. This equation can be solved variationally for $\bar{U}(R)$, at fixed values of R , by guessing the form the wave function, Ψ , and then minimizing $\bar{U} = \bar{T} + \bar{V}$ with respect to the parameters in Ψ .

Figs. 2.a, shows the results for linear, linear-plus-coulomb, and flux-bubble potential models with fixed-colour, using the old string-flip variational wave function,^e

$$\Psi_{\alpha\beta} = e^{-\sum_{\min\{Q\bar{q}\}} (\beta r_{Q\bar{q}})^\alpha}, \quad (7)$$

with parameters α^f and β . These models yields an average well depth of $\mathcal{O}(3) \text{ MeV}$, which is not enough to bind the Q_2 molecule. This suggests that

^dIn fact, it was this mini-laboratory which enabled us to developed a distributed minimization algorithm with $\tau_{\text{CPU}} \sim \mathcal{O}(p/m)$.⁴

^eFigs. 2.a, also shows results for the semi-relativistic Dirac-Hartree-Fock equation,

$$\sum_{\min\{Q\bar{q}\}} [\nabla_{\bar{q}}^2 + (m_q + \sigma r_{Q\bar{q}})^2] \Psi = U(R)^2 \Psi \quad (6)$$

(*cf.*, 5), for $m_q = 0$ and 330 MeV .

^fAnalysis was also done for fixed values of α , at 2 and 1.74, which shows a 1% variation in well depth, suggesting past assumptions about α are valid.

none of these models will lead to nuclear binding,⁹ or that perhaps something is wrong with the wave function itself.




Fig. 2: Binding energy as function of heavy-quark separation for the old (a) and new (b) wave functions with fixed-colour, and respective plots (c) and (d) for the flux-bubble model with moving-colour. Shown are the string-flip (linear and linear-plus-Coulomb) and flux-bubble (plus an $SU_c(3)$ composite model) models for both wave functions, and semi-relativistic plots for the old wave function with fixed-colour.

Although the Q_2 system is far removed from its H_2 , hydrogen molecule, cousin from a dynamical point of view and the motivations for achieving localization are quite different, it would seem plausible to use a similar *ansatz*.^{8,9}

$$\Psi_{\alpha,\beta} = e^{-\beta^\alpha(r_{\bar{q}_1 Q_1}^\alpha + r_{\bar{q}_2 Q_2}^\alpha)} + e^{-\beta^\alpha(r_{\bar{q}_2 Q_1}^\alpha + r_{\bar{q}_1 Q_2}^\alpha)}, \quad (8)$$

⁹We are not surprised that the flux-bubble model yields very little change, as the it was postulated that the colour-hyperfine interactions are more important.¹

where setting $\alpha = 1$ yields the H_2 case. Now, if $\bar{q}_1 Q_1$ and $\bar{q}_2 Q_2$ represent two separate mesons, say, then the first term represents the internal meson interactions while the second term represents the external meson interactions. Notice that the external interactions shut off as the separation, R , between the two heavy-quarks becomes large, which is the desired property.

Fig. 1.b, shows the results for linear, linear-plus-coulomb, and flux-bubble potential models with fixed-colour, using the new H_2 -like variational wave function.^h The contrast between the results for the old and new wave functions is quite dramatic! The new wave function gives a factor of $\mathcal{O}(27)$ increase in well depth; deep enough to bind a heavy-quark system with a reduced-mass greater than $\mathcal{O}(660)MeV$.⁴

Figs. 1.c and 1.d show the results for the flub-bubble models with moving-colour for the old and new wave function respectively. Fig. 1.c shows a linear core (with slope σ) with a repulsive barrier followed by an abrupt cut-off at $1.5fm$. This suggests that minimal energy state is a system consisting of two mesons, one containing heavy-quarks and the other containing light-antiquarks. Fig. 1.d shows potential with a very strong Coulomb-like (*i.e.*, $\alpha \sim \mathcal{O}(125.5)MeV fm$) interior, about $300MeV$ deep, with a Yukawa-like exterior. This suggests a very tightly bound Q_2 system, however, the Q_2 -well for the old wave function is much deeper, suggesting the aforementioned two-meson system is the minimal energy state. Of course this is not physical, and suggests that phenomenological models using $SU_c(2)$, with moving-colour, will yield unsatisfactory results.

Fig. 1.b, shows the results for an $SU_c(3)$ composite model, a “baryonic molecule,” which assumes the two heavy-quarks are actually a composite of two light-quarks (also, the orbiting antiquarks have been replaced with quarks). This model avoids dissociation into mesons, since a flux-tube can no longer be formed between the (composite) heavy-quarks, thus, restoring the potential to its generic (original) form.

It should be stressed that the Q_2 system is a toy-model, a pedagogical aid, used to gain better insight into the complex nature of nuclear interactions. Nuclear systems (at low temperature) consist of equal mass quarks, whereas, the Q_2 consists of unequal mass quarks. Therefore, what is applicable for Q_2 system may not translate very well to nuclear matter. That said, the results herein, suggest that the many-body wave function for nuclear matter should be of the form,

$$\Psi \sim \text{Perm} \left| e^{-(\beta r_{ij})^\alpha} \right| \prod_{\text{colour}} |\Phi(r_k)|. \quad (9)$$

^hAlso shown, are results for α fixed at 2 and 1.74.

which is a product of a totally symmetric correlation wave function times a totally antisymmetric Slater wave function. A simple test of this, *ansatz*, would be to consider Ψ in the context of a $q\bar{q}$ gas with fixed- $SU_c(2)$.

4 Conclusions

The Q_2 system has proven to be a very useful aid for trying to sort out the complexities of model building for nuclear matter. The details of the mechanics, from wave functions to dynamics to practical computing methods, of the flux-bubble model have been thoroughly investigated. However, it remains to be seen how applicable this work will be to nuclear matter; something that we hope to investigate in the near future.

In closing, it appears that the flux-bubble model may prove not only useful for modeling nuclear matter, but also useful for modeling mesonic molecules as well.

Acknowledgments

This work was supported by NSERC. The Monte Carlo calculations were performed on an 8 node DEC 5240 UNIX CPU farm, using Berkeley Sockets (TCP/IP).¹⁰ The computing facilities were provided by the Carleton University's Dept. of Physics, OPAL, CRPP, and THEORY. We would like to thank M.A. Doncheski, H. Blundell, M. Jones, and J.S. Wright for useful discussions.

References

1. M.M. Boyce & P.J.S. Watson, Nuc. Phys. **A580** (1994) 500; hep-ph/9312328.
2. C.J. Horowitz & J. Piekarewicz, Nucl. Phys. **A536** (1991) 669.
3. P.J.S. Watson, Nucl. Phys. **A494** (1989) 543.
4. M.M. Boyce, **String Inspired QCD and E₆ Models**, *Ph.D. thesis (1996)*, Carleton U., Dept. of Physics; hep-ph/9609433.
5. M.M. Boyce & P.J.S. Watson, *MRST '94 conf. proc.*, **ed.**, J.R. Cudell, K.R. Dienes, & B. Margolis, **pub.**, World Sci. (1994); hep-ph/9501397.
6. A.B. Migdal & P.J.S. Watson, Phys. Lett. **B252** (1990) 32.
7. J. Weinstein & N. Isgur, Phys. Rev. **D41** (1990) 2236, *ibid*, **D27** (1983) 588, *ibid*, Lett. (1982) 659.
8. Leonard I. Schiff, **Quantum Mechanics**, 2nd ed, McGraw Hill (1968).
9. B.H. Bransden & C.J. Joachain, **Physics of Atoms and Molecules**, Longman Group Limited (1983).
10. W.R. Stevens, **UNIX Network Programming**, Prentice Hall (1990).